

## GRB observations with H.E.S.S.

P. H. TAM<sup>(1)</sup>, S. J. WAGNER<sup>(1)</sup>, G. PÜHLHOFFER<sup>(1)</sup>

and THE H.E.S.S. COLLABORATION

<sup>(1)</sup> *Landessternwarte Heidelberg - Königstuhl, D-69117, Heidelberg, Germany*

**Summary.** — H.E.S.S. (High Energy Stereoscopic System), which is designed to detect TeV gamma-rays, is a system of four Imaging Atmospheric Cherenkov Telescopes situated in Namibia. The system has been shown to be very successful in detecting and observing galactic and extra-galactic TeV sources. In order to explore the highest energy end of GRB spectra, a GRB observing program has been established in the H.E.S.S. collaboration. Here we introduce our GRB observing program and report on its current status.

PACS 95.55.Ka – X- and gamma-ray telescopes and instrumentation.

PACS 98.70.Rz –  $\gamma$ -ray sources;  $\gamma$ -ray bursts.

### 1. – Very-high-energy emission from GRBs

The highest energy radiation from GRBs ever detected unambiguously by any instrument was a  $\sim 18$  GeV photon coming from GRB 940217 [1]. In the context of standard models of GRBs, photons with energies up to  $\sim 10$  TeV are expected. There could be an energy flux radiated in this largely-unexplored energy regime comparable to that radiated in keV-MeV or X-ray-to-radio energies. With the high sensitivity level of H.E.S.S. (for a source with integral flux  $\sim 1.4 \times 10^{-11}$  ph cm<sup>-2</sup>s<sup>-1</sup> above 1 TeV and spectral index 2.6, only a 2-hour observation with H.E.S.S. is needed for a  $5\text{-}\sigma$  detection ( $>10$  events)), we are capable to detect any signal comparable to that predicted in [2, 3].

### 2. – H.E.S.S. GRB observing program

A review of the system and observational highlights of H.E.S.S. can be found in [4].

Since early 2005, the Multi-wavelength Working Group in the H.E.S.S. collaboration has been responsible for possible GRB observations. Every month, a member (called the contact person) from the GRB sub-group is scheduled to be the coordinator between the team and the shift crew at the H.E.S.S. site for any possible GRB observations. A 24-hour-per-day monitoring is guaranteed.

We currently follow on-board GRB triggers distributed by *Swift*, as well as triggers from HETE II and INTEGRAL confirmed by ground-based analysis. Upon the reception

TABLE I. – 14 GRBs observed with H.E.S.S. from March 2003 to May 2006

| GRB     | Observation started after GRB onset | redshift   |
|---------|-------------------------------------|------------|
| 060526  | 4.7 h                               | 3.21 [5]   |
| 060505  | 19.4 h                              | 0.089 [6]  |
| 060403  | 13.6 h                              | –          |
| 050922C | 52 min                              | 2.199 [7]  |
| 050801  | 16 min                              | 1.45 [8]   |
| 050726  | 10.8 h                              | –          |
| 050607  | 14.8 h                              | –          |
| 050509C | 21 h                                | –          |
| 050209  | 20.2 h                              | –          |
| 041211  | 9.5 h                               | –          |
| 041006  | 10.4 h                              | 0.716 [9]  |
| 040425  | 26 h                                | –          |
| 030821  | 18 h                                | –          |
| 030329  | 11.5 d                              | 0.169 [10] |

of a new GRB notice (with good indications of being a true GRB) from the GRB Coordinates Network (GCN), we observe the reported burst position as soon as possible. The observation is limited to Z.A.  $\leq 45$  deg (to achieve a reasonably low energy threshold) and H.E.S.S. dark time which requires that the moon is down. The H.E.S.S. dark-time fraction is about 0.2. When a GRB observation is being performed, the GRB team members and the GRB contact person of that period are responsible for the ongoing observation and keep the shift crew updated of any new information about the GRB.

### 3. – Current status of the program

We have been observing GRBs since early 2003. At the beginning of 2005, a GRB coordination team was set up and since then our GRB observation program (see above) has been fully established. By May 2006, 14 GRB positions have been observed with H.E.S.S. (as shown in table I). The results will be published elsewhere [11].

\* \* \*

### REFERENCES

- [1] HURLEY K. ET AL., *Nature*, **372** (1994) 652.
- [2] ZHANG B. and MÉSZÁROS P., *Astrophys. J.*, **559** (2001) 110.
- [3] WANG X. Y. ET AL., *Astron. Astrophys.*, **439** (2005) 957.
- [4] HOFMANN W., *H.E.S.S. status*, in *Proc. Conf. Towards a Network of Atmospheric Cherenkov Detectors VII, Palaiseau, France* (2005), pp. 43-56.
- [5] BERGER E. and GLADDERS M., *GCN 5170*, (2006)
- [6] OFEK E. O., ET AL., *GCN 5123*, (2006)
- [7] D’ELIA V., ET AL., *GCN 4044*, (2005)
- [8] ROMING P., *this conference*, (2006)
- [9] PRICE P. A., ET AL., *GCN 2791*, (2004)
- [10] STANEK K. Z., ET AL., *Astrophys. J.*, **591** (2003) L17.
- [11] AHARONIAN F., ET AL., *in preparation*.